Published: March 25, 2019

REVIEW ARTICLE

Corresponding Author:

Daouda COULIBALY, MD, MPH

Address: BP V14, Abidjan,
National Institute of Public
Hygiene

Phone: +225 07 98 47 52

Email: daocoul@yahoo.fr

Declaration:

No conflict of interest was
reported.

Keywords: Outbreak, Ebola virus,
Côte d'Ivoire

How Côte d'Ivoire avoided the 2014 Ebola Virus Disease (EVD) outbreak.

D Coulibaly⁽¹⁾, EV Adjogoua⁽³⁾, D Chérif⁽¹⁾, Y Traoré^(1,2), A N'gattia⁽¹⁾, J Béné^(1,2), BM Dosso^(2,3), SN Dagnan^(1,2)

Author's affiliations:

¹. National Institute of Public Hygiene, Abidjan, Côte d'Ivoire

². Faculty of Medical Sciences - Université Félix Houphouët Boigny, Abidjan, Côte d'Ivoire

³. Pasteur Institute, Abidjan, Côte d'Ivoire

ABSTRACT

INTRODUCTION: The Ebola Virus Disease Epidemic (EVD), confirmed in March 2014 in Guinea, spread to Liberia, Sierra Leone, resulting in 28,646 cases over two years, including 11,323 deaths across ten countries. Côte d'Ivoire, which shares borders with Guinea and Liberia, due to high level of population movements, was considered a country at high risk of contamination. Despite its proximity to epidemic outbreaks, Côte d'Ivoire recorded no cases of Ebola virus disease. Faced with a threat posed by this dreadful disease, the Ivorian authorities put in place a series of measures to prevent the spread of the epidemic throughout the territory. The present work aims at describing a control strategy in order to increase knowledge on methods to prevent the cross-border spread of the epidemics.

PREVENTION STRATEGIES: The strategic axes were mainly coordination, surveillance, border health control, case management and community involvement.

RESULTS: A total of 92 alert cases including 10 suspects were reported for which the Ebola tests were negative. There were 66,880 phone calls recorded across the two hotlines. Following the confirmation of the epidemic on 21 March, 2014, four forward coordination posts were established in the border areas of Guinea and Liberia. These posts made it possible, with the involvement of the administrative authorities, opinion leaders, traditional and religious communities, to raise awareness and inform all the populations of the border localities about the methods of transmission and measures to prevent the disease, in particular hand washing, preventing the handling of corpses, hunting and the consumption of game.

CONCLUSION: Despite the measures taken to address the threat of Ebola, the risk of spreading the epidemic in Côte d'Ivoire existed. The strategies developed made it possible to prevent the cross-border spread of the epidemic in Côte d'Ivoire. The prevention measure relating to community involvement could be considered the fundamental factor that helped protect the Ivorian population against this dreadful disease. Certainly, there was a high risk of spreading the epidemic in Côte d'Ivoire; yet; not absolute.

Copyright:

© 2019 Science Publishing Group

This open access article is
distributed under the terms of the
[Creative Commons Attribution Non-
Commercial License](https://creativecommons.org/licenses/by-nc/4.0/).

INTRODUCTION

The Ebola Virus Disease (EVD) epidemic, the first of its kind in West Africa and confirmed in March 2014 in Guinea (1–3), spread to Liberia and Sierra Leone. Over two years, it resulted in 28,646 cases including 11,323 deaths across ten countries (4). In West Africa, cases were recorded in Nigeria, Mali and Senegal. Thanks to the efforts of the affected countries and the international community, the outbreaks were brought under control (5–9). The World Health Organization, then, declared the end of the public health emergency of international concern related to Ebola on 29 March, 2016 (10). In 2014, the epidemic was particularly rapid with more than 20,000 cases detected, including 10,000 in Guinea and Liberia, two countries bordering Côte d'Ivoire (4). Given the importance of cross-border population movements related to commercial and social activities, Côte d'Ivoire was considered a country at high risk of contamination (11–14). Indeed, the epidemic outbreaks were recorded in border localities namely Lola in Guinea and Nimba in Liberia (15). This risk of spread was also linked to historical factors, such as the discovery in 1994, in the Tai forest in western Côte d'Ivoire, of a non-lethal case of Ebola caused by the Ebolavirus species in Tai forest (3,12,13). Despite its proximity to epidemic outbreaks, Côte d'Ivoire was not affected by the epidemic. Faced with the threat posed by this dreadful disease, the Ivorian authorities put in place a series of measures to prevent the spread of the epidemic throughout the country. This piece of work aims at describing the control strategy in order to increase knowledge on the methods to prevent the cross-border spread of the epidemics.

PREVENTION STRATEGIES

The prevention strategies were coordination, surveillance, border health control, case management and community involvement.

Coordination

A monitoring committee chaired by the Prime Minister of the Republic of Côte d'Ivoire coordinated the control. This committee, which included ministers and ambassadors, met weekly to monitor the epidemic and adapt control measures. This committee set up four (04) forward coordination posts in the border regions with Guinea (Man, Odienné) and Liberia (Guiglo, San Pedro). These coordination posts were chaired by prefects of these regions and their task involved organising the control in accordance with the national guidelines drawn up by the monitoring committee. Following the confirmation of the epidemic on 21 March 2014 by WHO in Guinea (1), the forward coordination posts were established on 25 March 2014, four days later.

Epidemiological and virological surveillance.

The surveillance system put in place was an early warning system that concerned the whole country but was particularly focused on the border regions with Guinea and Liberia. This system was based on community surveillance, hospital surveillance and virological surveillance. Case definitions used were: alert case; anyone with high fever, sudden onset or bleeding or red diarrhoea or blood in the urine or sudden death; suspected case: Anyone suffering from a high fever ($T^{\circ} \geq 38^{\circ} \text{C}$) with or without bleeding (epistaxis, hematemesis, rectorragy) and associated with at least one of the following exposure factors: Staying in an area with an Ebola fever outbreak, having contact with a person infected or has died of Ebola, having contact with a dead or sick animal; confirmed case: suspected case confirmed by the laboratory (positive IgM serology, positive PCR or virus isolation), or having an epidemiological link with confirmed cases or an epidemic.

Thus, using the WHO-inspired case definition (16), suspected cases were reported to the National Institute of Public Hygiene (INHP), the coordinating structure for surveillance and response to epidemics. Suspected cases were reported immediately by health staff either electronically using a smartphone

through the MAGPI (advanced mobile data collection) application, or through other channels (phone, SMS, email). The community reported cases directly to health workers or through two hotlines (# 101 and # 143). These cases reported by the community were investigated by Rapid Response Teams. For any suspected case, a blood sample was taken and sent to the virology laboratory of Pasteur Institute of Côte d'Ivoire (IPCI) for analysis. Samples received at the laboratory were inactivated with AVL lysis buffer (Qiagen) followed by extraction of viral RNA using the QiaAmp viral mini kit (Qiagen). The amplification of the viral DNA was done by real-time RT-PCR with an ABI 7500 system thermocycler by primers and probes provided by the Institut Pasteur-Dakar and later with the Altona real star kit.

Border health control

Health controls concerned both the airport and maritime borders. Measures implemented at these entry points were: a systematic search for suspicious cases by taking the temperature using a thermal camera or an infrared thermometer, collecting information at the entry point using a sheet on a traveller's identity and exposure factors (staying in a risk area, having contact with an Ebola patient, handling a dead body, consuming game) and advising travellers on measures to prevent Ebola virus disease.

A special terminal was dedicated to flights from the three countries heavily affected by the Ebola virus disease (Guinea, Liberia, Sierra Leone). Thus, for travellers from these countries, any suspected case was referred by ambulance to the pandemics centre at Félix Houphouët Boigny International Airport and was the subject of a blood sample that was sent to the laboratory.

The land borders with Guinea and Liberia were closed in August 2014; however, with the possibility of a humanitarian and economic corridor that allowed the movement of people and the transfer of various products (medicines, equipment, food, goods) under certain conditions aimed at reducing the risk of spread of the disease in Côte d'Ivoire (disinfection of vehicles from affected countries, change of drivers of vehicles, temperature rise of drivers, quarantine for 21 days).

Case management

Five Ebola disease treatment centres were set up in the four border regions (Man, Guiglo, Odienné, San Pedro) and Abidjan. These centres were developed in accordance with the standards required for the management of highly contagious diseases (17,18). The staff of these centres were trained on the protocol of care of patients, biosafety, hospital hygiene, sample collection methods. Examination gloves and personal protective equipment were made available to all 82 health districts in the country to protect health personnel.

Community involvement

The community's commitment to implementing Ebola prevention measures was made through awareness, information and mobilisation campaigns. Mass awareness campaigns involving administrative authorities, opinion leaders, traditional and religious communities, NGOs, the private sector, the press were conducted throughout the country with a particular focus in the border regions of Guinea and Liberia. Village monitoring committees were set up in the four border regions to monitor the implementation of the preventive measures recommended by the administrative and health authorities. Three measures were the focus of attention: not eating bushmeat, not shaking hands, washing hands regularly with soap and water. To this end, handwashing devices and hydro-alcoholic solutions were placed in public places, in health institutions, at the airport, sea and land entry points. Large gatherings related to the organisation of common market days between border localities were also banned by the administrative authorities. Similarly, traditional healers should not claim to have Ebola treatment. Two hotlines were also made available to the population to give them information on the Ebola virus disease by a team of call center operators 24 hours a day.

Support from Partners

Several technical and financial partners as well as many NGOs assisted the Cote d'Ivoire authorities in managing this health threat related to EVD. For example, the World Health Organization, the US Centers for Diseases Control and Prevention, and Médecins Sans Frontières have contributed to the training of healthcare personnel in the areas of epidemiological surveillance, contact tracing, and case management at Ebola treatment centers.

RESULTS

Epidemiological surveillance

In West Africa, from 21 March 2014 to 27 March 2016, 28,639 cases of EVD were reported out of which 11,322 (CFR = 39.5%) died. The EVD cases reported in Guinea (3811 cases, 2,543 deaths), Liberia (10,675 cases, 4,809 deaths), Sierra Leone (14,124 cases, 3 956 deaths), Mali (8 cases, 6 deaths), Nigeria, 20 cases, 8 deaths), Senegal (1 case, 0 deaths). Among the most affected countries, the highest case fatality rates were observed in Guinea (66.7%), Liberia (45%) and Sierra Leone (28%) (Table 1). In Côte d'Ivoire, the epidemiological surveillance system recorded 92 alert cases during the year, including 7 deaths (CFR = 7.6%); 10 suspected cases not deceased throughout the country. None of these cases were confirmed by the laboratory as EVD (Table 2).

Table 1 : cases and deaths of Ebola Virus Disease, March 21, 2014 - March 27, 2016, West Africa

Country	Cases*	Deaths	CFR (%)
Guinea	3 811	2 543	66.7
Liberia	10 675	4 809	45.0
Sierra Léone	14 124	3 956	28.0
Mali	8	6	75.0
Nigeria	20	8	40.0
Senegal	1	0	0
Total	28 639	11 322	39.5

* : suspected, probable, confirmed

Table 2 : alert and suspect cases of Ebola Virus Disease, 2014, Côte d'Ivoire

Case definition	Cases	Deaths	CFR (%)
Alert case	92	7	7.6
Suspected case	10	0	0
Confirmed case	0		

Border health control

In each of the border localities of Côte d'Ivoire, Lola and Beyla in Guinea and Nimba in Liberia, the number of Ebola cases varied from 6 to 20 cases as of 31 December, 2014 (Figure 1). Out of 3,040 travellers from Guinea (1,763), Liberia (620) and Sierra Leone (657) respectively, five (05) suspected cases of Ebola among travellers from Guinea were reported through health checks carried out at Félix Houphouët Boigny International Airport in 2014. For these five cases, confirmation tests for Ebola were negative (Table 3).

Figure 1: geographical distribution of new and total confirmed cases in Guinea, Liberia, Mali and Sierra Leone, 2014.

Data for the past 21 days represent confirmed cases in Guinea (2014-12-20), Liberia (2014-12-18), Mali (2014-12-18) and Sierra Leone (2014-12-20).

LR: Liberia; SL: Sierra Leone; GI: Guinea; ML: Mali

Source: WHO

Table 3 : suspected cases detected at the airport among travelers from countries affected by the Ebola virus disease, 2014, Côte d'Ivoire.

Country	Number of travelers	Suspected case	Confirmed case
Guinea	1763	5	0
Liberia	620	0	0
Sierra Léone	657	0	0
TOTAL	3040	5	0

Use of the hotline by communities to learn about Ebola.

There were 66,880 calls recorded across the hotlines. The number of phone calls on the two hotlines to learn about the Ebola virus disease or to report cases gradually increased from March, date of confirmation of the epidemic in Guinea to 8,000 calls in August, and 14,000 in December 2014 after a decline in October with 6,000 calls (Figure 2).

Figure 2: Evolution of telephone calls on the green line and on the EVD, March to December 2014, Côte d'Ivoire.

DISCUSSION

The risk of the spread of the Ebola outbreak in Côte d'Ivoire remained a constant concern given the continuing epidemic in Guinea and Liberia. Indeed, the epidemic was beyond control in these countries and constituted a sub-regional and international threat (19–23). Risk assessment in the affected countries revealed factors of epidemic outbreak including community transmission linked to strong cultural and traditional practices and a community ignorance of the disease (24–26). The existence of an international airport and the sharing of two borders respectively with Guinea and Liberia further exposed Côte d'Ivoire to the disease. The speed and volume of air transport as well as cross-border trade were considered factors for the spread of epidemics (1,27–29). Coordination of the control that was the responsibility of the Prime Minister showed great importance given by the Ivorian authorities to the management of the health threat. This very high level of coordination enabled the mobilisation of populations and technical and financial partners for the implementation of prevention and preparedness interventions for the response. The effects of the epidemic in the affected countries on society (morbidity, mortality, unemployment), and the economy (tourism, air transport, fiscal resources) justified this real involvement of the authorities (30–34). The measures put in place in Côte d'Ivoire to cope with the epidemic began four days after the confirmation of the epidemic by setting up control coordination posts in the border regions of Guinea and Liberia. This promptness in the implementation of the interventions showed the authorities' willingness to prevent the spread of the disease or to detect it quickly. The epidemiological and virological surveillance system was strengthened to ensure the existence or absence of Ebola cases and to carry out necessary investigations. Between August and December 2014, the outbreak was very intense in the three affected countries raising fears of importing cases to Côte d'Ivoire (29). It was therefore important to have a surveillance system capable of checking warnings and confirming possible cases. The reported suspected cases were all negative for EVD. The detection of these cases showed the dynamic nature and the capacity of the surveillance system put in place. The closure of land borders with Guinea and Liberia was an exceptional measure that occurred when the epidemic was not controlled. This measure certainly helped to prevent the introduction of the disease into Côte d'Ivoire by reducing the access of patients and their contacts to the country, but it had limitations. Indeed, the borders are porous and cannot be closed over a long period because of political, diplomatic and economic consequences. This measure, which was not recommended by WHO, was also denounced by affected countries (11,12).

Among the Ebola threat management strategies, community involvement was prominent. The appropriation of the control by the communities was sought through several actions including information and awareness campaigns, the setting up of village monitoring committees, the effective involvement of religious and community leaders. These actions were implemented with determination and made it possible for the population to support the preventive measures. Thus, in places of worship and during funeral ceremonies people no longer shook hands (35–37). Traditional healers had also refrained from declaring their ability to treat Ebola (38). A story about the spread of the epidemic in Sierra Leone from a traditional healer justified the decision made by the Ivorian authorities (39).

CONCLUSION

Despite the measures taken to address the threat of Ebola, the risk of spreading the epidemic in Côte d'Ivoire existed. The strategies developed made it possible to prevent the cross-border spread of the epidemic in Côte d'Ivoire. The prevention measure relating to community involvement could be considered the fundamental factor that helped protect the Ivorian population against this dreadful disease. The high risk of the epidemic spreading to Côte d'Ivoire was not an absolute risk.

Acknowledgements: We express our sincere thanks to the Ebola Committee, national and international organizations and all those who have helped to prevent the spread of Ebola virus disease in Côte d'Ivoire. Special thanks to Dr. Raymonde GOUDOU who was Minister of Public Health.

REFERENCES

1. WHO | Six months after the Ebola outbreak was declared: Introduction [Internet]. WHO. [cité 7 janv 2019]. Disponible sur: <https://www.who.int/csr/disease/ebola/ebola-6-months/introduction/en/>
2. Guinea: Ebola epidemic declared [Internet]. 2014 [cité 7 janv 2019]. Disponible sur: <https://www.msf.org.uk/article/guinea-ebola-epidemic-declared>
3. Years of Ebola Virus Disease Outbreaks | 2014-2016 Outbreak West Africa | History | Ebola (Ebola Virus Disease) | CDC [Internet]. 2018 [cité 7 janv 2019]. Disponible sur: <https://www.cdc.gov/vhf/ebola/history/chronology.html>
4. WHO | Ebola situation reports: archive [Internet]. WHO. [cité 7 janv 2019]. Disponible sur: <http://www.who.int/csr/disease/ebola/situation-reports/archive/en/>
5. Fin de l'épidémie d'Ebola en Afrique de l'Ouest : le monde doit tirer les leçons pour la prochaine épidémie [Internet]. Médecins sans frontières. [cité 7 janv 2019]. Disponible sur: <https://www.msf.fr/communiqués-presse/fin-de-l-epidemie-d-ebola-en-afrique-de-l-ouest-le-monde-doit-tirer-les-lecons-pour-la-prochaine-epidemie>
6. Tuncer N, Mohanakumar C, Swanson S, Martcheva M. Efficacy of control measures in the control of Ebola, Liberia 2014-2015. *J Biol Dyn*. 2018;12(1):913-37.
7. Jalloh MF, Robinson SJ, Corker J, Li W, Irwin K, Barry AM, et al. Knowledge, Attitudes, and Practices Related to Ebola Virus Disease at the End of a National Epidemic - Guinea, August 2015. *MMWR Morb Mortal Wkly Rep*. 20 oct 2017;66(41):1109-15.
8. Dahl BA, Kinzer MH, Raghunathan PL, Christie A, De Cock KM, Mahoney F, et al. CDC's Response to the 2014-2016 Ebola Epidemic - Guinea, Liberia, and Sierra Leone. *MMWR Suppl*. 8 juill 2016;65(3):12-20.
9. Al TGN et. Ebola and Its Control in Liberia, 2014–2015 - Volume 22, Number 2—February 2016 - Emerging Infectious Diseases journal - CDC. [cité 7 janv 2019]; Disponible sur: https://wwwnc.cdc.gov/eid/article/22/2/15-1456_article
10. Ebola is no longer a public health emergency [Internet]. [cité 7 janv 2019]. Disponible sur: <https://www.who.int/news-room/detail/29-03-2016-statement-on-the-9th-meeting-of-the-ihf-emergency-committee-regarding-the-ebola-outbreak-in-west-africa>
11. WHO | Statement on the 1st meeting of the IHR Emergency Committee on the 2014 Ebola outbreak in West Africa [Internet]. WHO. [cité 7 janv 2019]. Disponible sur: <https://www.who.int/mediacentre/news/statements/2014/ebola-20140808/en/>
12. WHO | Statement on the 3rd meeting of the IHR Emergency Committee regarding the 2014 Ebola outbreak in West Africa [Internet]. WHO. [cité 7 janv 2019]. Disponible sur: <https://www.who.int/mediacentre/news/statements/2014/ebola-3rd-ihf-meeting/en/>
13. EVD_WestAfrica_WHO_RiskAssessment_20140624.pdf [Internet]. [cité 7 janv 2019]. Disponible sur: https://www.who.int/csr/disease/ebola/EVD_WestAfrica_WHO_RiskAssessment_20140624.pdf?ua=
14. Bogoch II, Creatore MI, Cetron MS, Brownstein JS, Pesik N, Miniota J, et al. Assessment of the potential for international dissemination of Ebola virus via commercial air travel during the 2014 west African outbreak. *The Lancet*. 3 janv 2015;385(9962):29-35.

15. WHO | Ebola maps [Internet]. WHO. [cité 7 janv 2019]. Disponible sur: <http://www.who.int/csr/disease/ebola/maps/en/>
16. WHO | Case definition recommendations for Ebola or Marburg virus diseases [Internet]. WHO. [cité 7 janv 2019]. Disponible sur: <https://www.who.int/csr/resources/publications/ebola/case-definition/en/>
17. WHO | Clinical management of patients with viral haemorrhagic fever [Internet]. WHO. [cité 7 janv 2019]. Disponible sur: <http://www.who.int/csr/resources/publications/clinical-management-patients/en/>
18. En savoir plus sur l'Ebola [Internet]. Médecins sans frontières. [cité 7 janv 2019]. Disponible sur: <https://www.msf.fr/actualites/en-savoir-plus-sur-l-ebola>
19. Loungou S. L'épidémie d'Ebola en Afrique de l'Ouest. Une mise en perspective des répercussions démo-géographiques, politiques et économiques. L'Espace Polit Rev En Ligne Géographie Polit Géopolitique [Internet]. 22 juill 2015 [cité 7 janv 2019];(26). Disponible sur: <http://journals.openedition.org/espacepolitique/3467>
20. WHO | Barriers to rapid containment of the Ebola outbreak [Internet]. WHO. [cité 7 janv 2019]. Disponible sur: <https://www.who.int/csr/disease/ebola/overview-august-2014/en/>
21. WHO | Ebola in West Africa: heading for catastrophe? [Internet]. WHO. [cité 7 janv 2019]. Disponible sur: <https://www.who.int/csr/disease/ebola/ebola-6-months/west-africa/en/>
22. Ebola Virus Disease in West Africa — The First 9 Months of the Epidemic and Forward Projections. N Engl J Med. 16 oct 2014;371(16):1481-95.
23. msf_ebolareport_fr-def.pdf [Internet]. [cité 7 janv 2019]. Disponible sur: https://www.msf.fr/sites/default/files/2018-03/msf_ebolareport_fr-def.pdf
24. WHO | Factors that contributed to undetected spread of the Ebola virus and impeded rapid containment [Internet]. WHO. [cité 7 janv 2019]. Disponible sur: <http://www.who.int/entity/csr/disease/ebola/one-year-report/factors/en/index.html>
25. Rapp C. [Ebola virus epidemic in West Africa: facts and prospects]. Med Sante Trop. sept 2014;24(3):229-31.
26. Carrión Martín AI, Derrough T, Honomou P, Kolie N, Diallo B, Koné M, et al. Social and cultural factors behind community resistance during an Ebola outbreak in a village of the Guinean Forest region, February 2015: a field experience. Int Health. 1 mai 2016;8(3):227-9.
27. Undurraga EA, Carias C, Meltzer MI, Kahn EB. Potential for broad-scale transmission of Ebola virus disease during the West Africa crisis: lessons for the Global Health security agenda. Infect Dis Poverty. 1 déc 2017;6(1):159.
28. Cowling BJ, Yu H. Ebola: worldwide dissemination risk and response priorities. Lancet Lond Engl. 3 janv 2015;385(9962):7-9.
29. Gomes MFC, Pastore y Piontti A, Rossi L, Chao D, Longini I, Halloran ME, et al. Assessing the International Spreading Risk Associated with the 2014 West African Ebola Outbreak. PLoS Curr [Internet]. 2 sept 2014 [cité 7 janv 2019];6. Disponible sur: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4169359/>

30. Bank TW. The economic impact of the 2014 Ebola epidemic : short and medium term estimates for West Africa [Internet]. The World Bank; 2014 oct [cité 7 janv 2019] p. 1-71. Report No.: 91219. Disponible sur: <http://documents.worldbank.org/curated/en/524521468141287875/The-economic-impact-of-the-2014-Ebola-epidemic-short-and-medium-term-estimates-for-West-Africa>
31. Cost of the Ebola Epidemic | 2014-2016 Outbreak West Africa | History | Ebola (Ebola Virus Disease) | CDC [Internet]. 2018 [cité 7 janv 2019]. Disponible sur: <https://www.cdc.gov/vhf/ebola/history/2014-2016-outbreak/cost-of-ebola.html>
32. Kirigia JM, Masiye F, Kirigia DG, Akweongo P. Indirect costs associated with deaths from the Ebola virus disease in West Africa. Infect Dis Poverty [Internet]. 29 oct 2015 [cité 7 janv 2019];4. Disponible sur: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4625462/>
33. Huber C, Finelli L, Stevens W. The Economic and Social Burden of the 2014 Ebola Outbreak in West Africa. J Infect Dis. 22 nov 2018;218(suppl_5):S698-704.
34. Ebola: Most African Countries Avoid Major Economic Loss but Impact on Guinea, Liberia, Sierra Leone Remains Crippling [Internet]. World Bank. [cité 7 janv 2019]. Disponible sur: <http://www.worldbank.org/en/news/press-release/2015/01/20/ebola-most-african-countries-avoid-major-economic-loss-but-impact-on-guinea-liberia-sierra-leone-remains-crippling>
35. Ebola : Les TIC pour combattre le fléau en Côte d'Ivoire [Internet]. [cité 7 janv 2019]. Disponible sur: <http://www.sciencepresse.qc.ca/blogue/2014/09/16/ebola-tic-pour-combattre-fleau-cote-divoire>
36. Côte d'Ivoire: la lutte contre Ebola s'intensifie sur Internet - RFI [Internet]. RFI Afrique. [cité 7 janv 2019]. Disponible sur: <http://www.rfi.fr/afrique/20140822-cote-ivoire-lutte-mousser-contre-ebola-intensifie-internet-reseaux-sociaux>
37. La Côte d'Ivoire en état d'alerte face à la menace de l'épidémie d'Ebola. 18 août 2014 [cité 7 janv 2019]; Disponible sur: https://www.lemonde.fr/planete/article/2014/08/18/la-cote-d-ivoire-en-etat-d-alerte-face-a-la-menace-de-l-epidemie-d-ebola_4472763_3244.html
38. Ebola : la Côte d'Ivoire met en garde les guérisseurs traditionnels [Internet]. [cité 7 janv 2019]. Disponible sur: https://www.rti.ci/actualite_article.php?categorie=societe&id=8105&titre=ebola-la-cote-deivoire-met-en-garde-les-guerisseurs-traditionnels&page=1
39. WHO | Sierra Leone: a traditional healer and a funeral [Internet]. WHO. [cité 7 janv 2019]. Disponible sur: <https://www.who.int/csr/disease/ebola/ebola-6-months/sierra-leone/en/>